arXiv:0811.1060v1 [math.RA] 6 Nov 2008

SUBNORMAL SUBALGEBRAS OF LEIBNIZ ALGEBRAS

DONALD W. BARNES

Abstract. Zassenhaus has proved that if U is a subnormal subalgebra of a finite-dimensional Lie algebra L and V is a finite-dimensional irreducible L-module, then all U-module composition factors of V are isomorphic. Schenkman has proved that if U is a subnormal subalgebra of a finite-dimensional Lie algebra L, then the nilpotent residual of U is an ideal of L. These useful results generalise to Leibniz algebras.

1. Introduction

Zassenhaus has proved [5, Lemma 1] that if U is a subnormal subalgebra of a finite-dimensional Lie algebra L and V is a finite-dimensional irreducible L-module, then all U-module composition factors of V are isomorphic. Zassenhaus's proof does not extends to Leibniz algebras. However, the result for Leibniz algebras is easily deduced from the Lie algebra result. Schenkman [4, Theorem 3] has proved that, for a subnormal subalgebra U of a Lie algebra L, the nilpotent residual $U_{\mathfrak{N}} = U^{\omega} = \bigcap_r U^r$ of U is an ideal of L. I show that this holds for Leibniz algebras. Schenkman's proof shows for a subnormal subalgebra U of a Leibniz algebra L, that $U_{\mathfrak{N}}$ is a right ideal of L, but a new argument is required to show that it is also a left ideal.

In the following, L is a finite-dimensional (left) Leibniz algebra over an arbitrary field, that is, an algebra whose left multiplication operators $\lambda_a\colon L \to L$ given by $\lambda_a(x) = ax$ for $a, x \in L$, are derivations. Thus $a(xy) = (ax)y + x(ay)$. The basic properties of Leibniz algebras and their bimodules may be found in Ayupov and Omirov [1], Patsourakos [3] or Barnes [2].

The left centre of a Leibniz algebra L is $\mathcal{Z}^l(L) = \{x \in L \mid xa = 0 \text{ for all } a \in L\}$. It is a two-sided ideal of L. For all $x \in L$, $x^2 \in \mathcal{Z}^l(L)$ and so $L/\mathcal{Z}^l(L)$ is a Lie algebra. If V is an L-bimodule, the centraliser of V in L is the two-sided ideal $\mathcal{C}_L(V) = \{x \in L \mid xv = vx = 0 \text{ for all } v \in V\}$. I write $U \lhd_{\mathrm{r}} L$ for "U is a right ideal of L" and $U \lhd L$ for "U is an ideal of L". A subalgebra U of L is called subnormal, written $U \lhd\!\lhd L$, if there exists a chain of subalgebras $U_0 = L > U_1 > \cdots > U_n = U$ with each $U_i \lhd U_{i-1}$.

2. The proofs

Lemma 2.1. *Let L be a finite-dimensional Leibniz algebra and let V be a finite-dimensional irreducible L-bimodule. Then $L/\mathcal{C}_L(V)$ is a Lie algebra and either $VL = 0$ or $vx = -xv$ for all $x \in L$ and $v \in V$.*

Proof. Without loss of generality, we may suppose $\mathcal{C}_L(V) = 0$. Form the split extension X of V by L and let $K = \mathcal{Z}^l(X)$. Since V is a minimal ideal of X, either $K \geq V$ or $K \cap V = 0$. As K is an abelian ideal, in either case, we have $K \leq \mathcal{C}_X(V)$. For all $x \in L$, $x^2 \in K$, so $x^2 = 0$. Thus L is a Lie algebra. As $X/V \simeq L$ is a Lie

---

2000 *Mathematics Subject Classification.* Primary 17A32.

Key words and phrases. Leibniz algebras, subnormal.

This work was done while the author was an Honorary Associate of the School of Mathematics and Statistics, University of Sydney.

algebra, $K \leq V$. Thus $K = V$ or $K = 0$. If $K = V$, then $VX = 0$. If $K = 0$, then X is a Lie algebra, so $vx = -xv$ for all $x \in L$ and $v \in V$. ☐

Theorem 2.2. *Let L be a finite-dimensional Leibniz algebra and let $U \lhd\!\lhd L$. Let V be a finite-dimensional irreducible L-bimodule. Then all U-bimodule composition factors of V are isomorphic.*

Proof. We can work with $L/\mathcal{C}_L(V)$, so we may suppose that $\mathcal{C}_L(V) = 0$. By Lemma 2.1, L is a Lie algebra and either $VL = 0$ or $vx = -xv$ for all $x \in L$ and $v \in V$. In either case, any left U-submodule is a U-sub-bimodule. All left U-module composition factors of V are isomorphic by Zassenhaus [5, Lemma 1] and the result follows. ☐

I write λ_U for the operator on the set of subspaces of L given by $\lambda_U V = UV$, the space of products uv of $u \in U$ and $v \in V$.

Lemma 2.3. *For any subspace $V \subseteq L$, $U^k V \subseteq \lambda_U^k V$.*

Proof. We use induction over k. The result holds trivially for $k = 1$. We have

$$U^{k+1}V = (UU^k)V \subseteq U(U^kV) + U^k(UV) \subseteq U(\lambda_U^k V) + \lambda_U^k(UV) = \lambda_U^{k+1}V,$$

by induction. ☐

Corollary 2.4. *Suppose $U \lhd\!\lhd L$. Then $U_{\mathfrak{N}} \lhd_{\mathrm{r}} L$.*

Proof. Suppose $U_0 = L > U_1 > \cdots > U_r = U$ is a chain of subalgebras with each $U_i \lhd U_{i-1}$. For some s we have $U_{\mathfrak{N}} = U^s$. We have

$$U_{\mathfrak{N}}L = U^{r+s}L \subseteq \lambda_U^{r+s}L \subseteq U^s = U_{\mathfrak{N}}$$

and $U_{\mathfrak{N}} \lhd_{\mathrm{r}} L$ as asserted. ☐

Theorem 2.5. *Suppose $U \lhd\!\lhd L$. Then $U_{\mathfrak{N}} \lhd L$.*

Proof. Suppose $U_0 = L > U_1 > \cdots > U_r = U$ is a chain of subalgebras with each $U_i \lhd U_{i-1}$ and that $U_{\mathfrak{N}} = U^s$. We use induction over r to prove that $U_{\mathfrak{N}}$ is a left ideal. This holds trivially for $r = 0$. Suppose $r > 0$. We use induction over t to show that $LU_{\mathfrak{N}} \subseteq \lambda_U^t L + U_{\mathfrak{N}}$. This holds trivially for $t = 0$. We have

$$LU_{\mathfrak{N}} = L(UU^s) \subseteq (LU)U^s + U(LU^s) \subseteq U_1U_{\mathfrak{N}} + U(\lambda_U^{t-1}L + U_{\mathfrak{N}}) \subseteq U_{\mathfrak{N}} + \lambda_U^t L$$

as $U_1U_{\mathfrak{N}} \subseteq U_{\mathfrak{N}}$ by induction over r. For $t = r + s$, $\lambda_U^t L \subseteq U_{\mathfrak{N}}$. ☐

References

[1] Sh. A. Ayupov and B. A. Omirov, *On Leibniz algebras,* Algebra and Operators Theory, Proceedings of the Colloquium in Tashkent, Kluwer, (1998) 1–13.

[2] D. W. Barnes, *Engel subalgebras of Leibniz algebras,* arXiv:0810.2849.

[3] A. Patsourakos, *On nilpotent properties of Leibniz algebras* Comm. in Algebra **35** (2007) 3828–3834.

[4] E. Schenkman, *A theory of subinvariant Lie algebras,* Amer. J. Math. **73** (1951), 453–474.

[5] H. Zassenhaus, *On trace bilinear forms on Lie-algebras,* Proc. Glasgow Math. Assoc. **4** (1959), 62–72.

1 Little Wonga Rd, Cremorne NSW 2090 Australia
E-mail address: `donwb@iprimus.com.au`